§ 866.3290

be transmitted to humans through handling of infected animals, animal products, or by the bites of fleas and ticks. The disease takes on several forms depending upon the site of infection, such as skin lesions, lymph node enlargements, or pulmonary infection.

(b) Classification. Class II (special controls). The device is exempt from the premarket notification procedures in subpart E of part 807 of this chapter subject to §866.9.

[47 FR 50823, Nov. 9, 1982, as amended at 63 FR 59226, Nov. 3, 1998]

§866.3290 Gonococcal antibody test (GAT).

(a) Identification. A gonococcal antibody test (GAT) is an in vitro device that consists of the reagents intended to identify by immunochemical techniques, such as latex agglutination, indirect fluorescent antibody, radioimmunoassav. antibodies to Neisseria gonorrhoeae in sera of asymptomatic females at low risk of infection. Identification of antibodies with this device may indicate past or present infection of the patient with Neisseria gonorrhoeae.

(b) Classification. Class III (premarket approval) (transitional device).

(c) Date PMA or notice of completion of a PDP is required. As of May 28, 1976, an approval under section 515 of the act is required before this device may be commercially distributed. See §866.3.

[47 FR 50823, Nov. 9, 1982, as amended at 52 FR 17734, May 11, 1987]

§866.3300 Haemophilus spp. serological reagents.

(a) Identification, Haemophilus spp. serological reagents are devices that consist of antigens and antisera, including antisera conjugated with a fluorescent dye, that are used in serological tests to identify Haemophilus spp. directly from clinical specimens or tissue culture isolates derived from clinical specimens. The identification aids in the diagnosis of diseases caused by bacgenus belonging to Haemophilus and provides epidemiological information on diseases cause by these microorganisms. Diseases most often caused by Haemophilus spp. include pneumonia, pharyngitis, sinusitis, vaginitis, chancroid venereal disease, and a contagious form of conjunctivitis (inflammation of eyelid membranes).

(b) Classification. Class II (special controls). The device is exempt from the premarket notification procedures in subpart E of part 807 of this chapter subject to §866.9.

[47 FR 50823, Nov. 9, 1982, as amended at 63 FR 59226, Nov. 3, 1998]

§866.3305 Herpes simplex virus serological reagents.

(a) Identification. Herpes simplex virus serological reagents are devices that consist of antigens and antisera used in various serological tests to identify antibodies to herpes simplex virus in serum. Additionally, some of the reagents consist of herpes simplex virus antisera conjugated with a fluorescent dye (immunofluorescent reagents) used to identify herpes simplex virus directly from clinical specimens or tissue culture isolates derived from clinical specimens. The identification aids in the diagnosis of diseases caused by herpes simplex viruses and provides epidemiological information on these diseases. Herpes simplex viral infections range from common and mild lesions of the skin and mucous membranes to a severe form of encephalitis (inflammation of the brain). Neonatal herpes virus infections range from an mild infection to a severe generalized disease with a fatal outcome.

(b) Classification. Class III (premarket approval).

(c) Date PMA or notice of completion of a PDP is required. No effective date has been established of the requirement for premarket approval. See §866.3.

 $[47\ FR\ 50823,\ Nov.\ 9,\ 1982,\ as\ amended\ at\ 52\ FR\ 17734,\ May\ 11,\ 1987]$

§866.3320 Histoplasma capsulatum serological reagents.

(a) Identification. Histoplasma capsulatum serological reagents are devices that consist of antigens and antisera used in serological tests to identify antibodies to Histoplasma capsulatum in serum. Additionally, some of these reagents consist of Histoplasma capsulatum antisera conjugated with a fluorescent dye (immunofluorescent reagents) used to identify Histoplasma capsulatum from

Food and Drug Administration, HHS

clinical specimens or cultured isolates derived from clinical specimens. The identification aids in the diagnosis of histoplasmosis caused by this fungus belonging to the genus *Histoplasma* and provides epidemiological information on the diseases caused by this fungus. Histoplasmosis usually is a mild and often asymptomatic respiratory infection, but in a small number of infected individuals the lesions may spread to practically all tissues and organs.

(b) Classification. Class II (special controls). The device is exempt from the premarket notification procedures in subpart E of part 807 of this chapter subject to §866.9.

[47 FR 50823, Nov. 9, 1982, as amended at 63 FR 59227, Nov. 3, 1998]

§866.3330 Influenza virus serological reagents.

(a) Identification. Influenza virus serological reagents are devices that consist of antigens and antisera used in serological tests to identify antibodies to influenza in serum. The identification aids in the diagnosis of influenza (flu) and provides epidemiological information on influenza. Influenza is an acute respiratory tract disease, which is often epidemic.

(b) Classification. Class I (general controls). The device is exempt from the premarket notification procedures in subpart E of part 807 of this chapter subject to the limitations in §866.9.

 $[47\ FR\ 50823,\ Nov.\ 9,\ 1982,\ as\ amended\ at\ 54\ FR\ 25047,\ June\ 12,\ 1989;\ 66\ FR\ 38792,\ July\ 25,\ 2001]$

\$866.3340 Klebsiella spp. serological reagents.

(a) Identification. Klebsiella spp. serological reagents are devices that consist of antigens and antisera, including antisera conjugated with a fluorescent dye (immunofluorescent reagents), that are used in serological tests to identify Klebsiella spp. from cultured isolates derived from clinical specimens. The identification aids in the diagnosis of diseases caused by bacteria belonging to the genus Klebsiella and provides epidemiological information on these diseases. These organisms can cause serious urinary tract and pulmonary infections, particularly in hospitalized patients.

(b) Classification. Class I (general controls). The device is exempt from the premarket notification procedures in subpart E of part 807 of this chapter subject to the limitations in §866.9.

[47 FR 50823, Nov. 9, 1982, as amended at 54 FR 25047, June 12, 1989; 66 FR 38792, July 25, 2001]

§866.3350 *Leptospira* spp. serological reagents.

(a) Identification. Leptospira spp. serological reagents are devices that consist of antigens and antisera used in serological tests to identify antibodies to Leptospira spp. in serum or identify Leptospira spp. from cultured isolates derived from clinical specimens. Additionally, some of these antisera are conjugated with a fluorescent dve (immunofluorescent reagents) and used to identify Leptospira spp. directly from clinical specimens. The identification aids in the diagnosis of leptospirosis caused by bacteria belonging to the genus Leptospira and provides epidemiological information on this disease. Leptospira infections range from mild fever-producing illnesses to severe liver and kidney involvement producing hemorrhage and dysfunction of these organs.

(b) Classification. Class II (special controls). The device is exempt from the premarket notification procedures in subpart E of part 807 of this chapter subject to §866.9.

[47 FR 50823, Nov. 9, 1982, as amended at 63 FR 59227, Nov. 3, 1998]

§866.3355 *Listeria* spp. serological reagents.

(a) Identification. Listeria spp. serological reagents are devices that consist of antigens and antisera used in serological tests to identify Listeria spp. from cultured isolates derived from clinical specimens. Additionally, some of these reagents consist of *Listeria* spp. antisera conjugated with a fluorescent dye (immunofluorescent reagents) used to identify Listeria spp. directly from clinical specimens. The identification aids in the diagnosis of listeriosis, a disease caused by bacteria belonging to the genus Listeria, and provides epidemiological information on diseases caused by these microorganisms. Listeria monocytogenes, the most common